



Fig. 1. Comparison of the reactivity of LK-1 MoAb with standard anti-CD43 MoAb (MEM-59) on cells of the CEM line and on CD43-deficient cells of the CEM line detected by flow cytometry. Line A: reactivity of LK-1 and standard anti-CD43 (MEM-59) MoAbs with CEM cells. A(1) Negative control double staining IgG1 FITC and IgG2 PE, A(2) comparison of the reactivity of LK-1 FITC and MEM-59 PE, double staining, A(3) comparison of the reactivity of MEM-59 FITC and LK-1 PE MoAbs, double staining. Line B: reactivity of LK-1 and standard anti-CD43 (MEM-59) MoAbs with CD43-deficient CEM cells. B(1) Negative control, double staining IgG1 FITC and IgG2 PE, B(2) comparison of the reactivity of MEM-59 FITC and LK-1 PE, double staining, B(3) comparison of the reactivity of LK-1 FITC and MEM-59 PE, double staining.

References

- Galfré, L., Milstein, C. (1981) Preparation of monoclonal antibodies: strategies and procedures. *Methods Enzymol.* **73**, 3-46.
- Hořejší, V., Stockinger, H. (1997) CD43 workshop panel report. In: *Leukocyte Typing VI: White Cell Differentiation Antigens*, eds. Kishimoto, T., Kikutani, H., von dem Borne, A., Goyert, S. M., Mason, D. Y., Miyasaka, M., Moretta, L., Okumura, K., Shaw, S., Springer, T. A., Sugamura, K., Zola, H., pp. 494-497, Garland Publishing, Inc., New York.
- Koubek, K. (1989) Monoclonal antibody LK-1 reactivity with human haematopoietic cells. *Sborník lék.* **91**, 242-248. (in Czech)
- Koubek, K., Stary, J., Kumberová, A., Klamová, H., Filipec, M. (1999) Occurrence of cytokine receptors on different lymphoid leukaemic cells. *Eur. J. Haemat.* **63**, 1-10.
- Manjunath, N., Johnson, R. S., Staunton, D. E., Pasqualini, R., Ardman, B. (1993) Targeted disruption of CD43 gene enhances T lymphocyte adhesion. *J. Immunol.* **151**, 1528-1534.
- Ostberg, J. R., Barth, R. K., Frelinger, J. G. (1998) The Roman god Janus: a paradigm for the function of CD43. *Immunol. Today* **12**, 546-550.
- Pallant, V., Eskenazi, A., Mattei, M. G., Fournier, R. E. K., Carlsson, S. R., Fukuda, M., Frelinger, J. G. (1989) Characterization of cDNAs encoding human leukosialin and localization of the leukosialin gene to chromosome 16. *Proc. Natl. Acad. Sci. USA* **86**, 1328-1332.
- Schmid, K., Hediger, M. A., Brossmer, R., Collins, J. H., Haupt, H., Marti, T., Offner, G. D., Schaller, J., Takagaki, K., Walsh, M. T. (1992) Amino acid sequence of human plasma galactoglycoprotein: identity with the extracellular region of CD43 (sialophorin). *Proc. Natl. Acad. Sci. USA* **89**, 663-667.
- Seveau, S., Keller, H., Maxfield, F. R., Piller, F., Halbwasch-Mecarelli, L. (2000) Neutrophil polarity and locomotion are associated with surface redistribution of leukosialin (CD43), an antiadhesive membrane molecule. *Blood* **95**, 2462-2470.
- Shaw, S. (1997) CD Guide. In: *Leukocyte Typing VI: White Cell Differentiation Antigens*, eds. Kishimoto, T., Kikutani, H., von dem Borne, A., Goyert, S. M., Mason, D. Y., Miyasaka, M., Moretta, L., Okumura, K., Shaw, S., Springer, T. A., Sugamura, K., Zola, H., pp. 1109-1220, Garland Publishing, Inc., New York.