

- myeloid leukemia by fluorescence in situ hybridization: comparison with conventional cytogenetics and implications for minimal residual disease detection. *Hum. Pathol.* **28**, 154-159.
- Heissig, B., Schultheis, B., Hochhaus, A., Pasternak, G., Hehlmann, R. (1997) A significant proportion of bcr-abl positive hematopoietic progenitors is present after long-term culture of peripheral blood mononuclear cells from patients on interferon- α therapy. *J. Mol. Med.* **75**, B247. (Abstract)
- Hörner, S., Pasternak, G., Hehlmann, R. (1997) A statistically significant sex difference in the number of colony-forming cells from human peripheral blood. *Ann. Hematol.* **74**, 259-263.
- Lichter, P., Cremer, T. (1992) Chromosome analysis by non-isotopic in situ hybridization. In: *Human Cytogenetics*, eds. Rooney, D., Czepulkowski, B., pp. 157, Oxford, University Press, New York.
- Pasternak, G., Pasternak, L. (1994) Persistence of bcr-abl mRNA-expressing cells in long term cultures established from chronic myeloid leukemic bone marrow or blood. *Ann. Hematol.* **68**, 9-14.
- Reiter, A., Marley, S. B., Hochhaus, A., Sohal, J., Raanani, P., Hehlmann, R., Gordon, M. Y., Goldman, J. M., Cross, N. C. P. (1998) BCR-ABL-positive progenitors in chronic myeloid leukaemia patients in complete cytogenetic remission after treatment with interferon- α . *Br. J. Haematol.* **102**, 1271-1278.
- Schultheis, B., Heissig, B., Sick, C., Hochhaus, A., Pasternak, G., Hehlmann, R. (1997) Long-term cultured colony-forming cells from peripheral blood from CML patients under interferon-alpha or hydroxyurea therapy show differences in BCR-ABL positivity. *Blood* **90**, 282b. (Abstract)
- Tkachuk, D., Westbrook, C., Andreeff, M., Donlon, T., Cleary, M., Suryanarayan, K., Homge, M., Redner, A., Gray, J., Pinkel, D. (1990) Detection of bcr-abl fusion in chronic myelogenous leukemia by in situ hybridization. *Science* **250**, 559-562.
- Udomsakdi, C., Eaves, C., Swolin, B., Reid, D., Barnett, M., Eaves, A. (1992) Rapid decline of chronic myeloid leukemic cells in long-term culture due to a defect at the leukemic stem cell level. *Proc. Natl. Acad. Sci. USA* **89**, 6192-6196.
- Verfaillie, C., Bhatia, R., Miller, W., Mortari, F., Roy, V., Burger, S., McCullough, J., Stieglbauer, K., Dewald, G., Heimfeld, S., Miller, J., McGlave, P. (1992a) BCR/ABL-negative primitive progenitors suitable for transplantation can be selected from the marrow of most early-chronic phase but not accelerated-phase chronic myelogenous leukemia patients. *Blood* **87**, 4770-4779.
- Verfaillie, C., Miller, W., Boylan, K., McGlave, P. (1992b) Selection of benign primitive hematopoietic progenitors in chronic myelogenous leukemia on the basis of HLA-DR antigen expression. *Blood* **79**, 1003-1010.