

- Křen, V. (1975) Genetics of the polydactyly-luxate syndrome in the Norway rat, *Rattus norvegicus*. *Acta Univ. Carol. Med. (Monogr.)* **68**, 1-104.
- Křen, V., Křenová, D., Pravenec, M., Zdobinská, M. (1995) Chromosome 8 congenic strains: tools for genetic analysis of limb malformation, plasma triglycerides and blood pressure in the rat. *Folia Biol. (Praha)* **41**, 284-293.
- Křen, V., Bílá, V., Kašpárek, R., Křenová, D., Pravenec, M., Rapp, K. (1996) Recombinant inbred and congenic strains of the rat for genetic analysis of limb morphogenesis. *Folia Biol. (Praha)* **42**, 159-166.
- Křen, V., Křenová, D., Šimáková, M., Musilová, A., Zídek, V., Pravenec, M. (2000) SHR.BN-congenic strains for genetic analysis of multifactorially determined traits. *Folia Biol. (Praha)* **46**, 25-30.
- Kwasigroch, T. E., Kochhar, D. M. (1980) Production of congenital limb defects with retinoic acid: phenomenological evidence of progressive differentiation during limb morphogenesis. *Anat. Embryol. (Berl.)* **161**, 105-113.
- Lau, W. F., Tertinegg, I., Heersche, J. N. (1993) Effects of retinoic acid on cartilage differentiation in a chondrogenic cell line. *Teratology* **47**, 555-563.
- Lee, K. K., Li, F. C., Yung, W. T., Kung, J. L., Ng, J. L., Cheah, K. S. (1994) Influence of digits, ectoderm, and retinoic acid on chondrogenesis by mouse interdigital mesoderm in culture. *Dev. Dyn.* **201**, 297-309.
- Lohnes, D., Mark, M., Mendelsohn, C., Dolle, P., Dierich, A., Gorry, P., Gansmuller, A., Chambon, P. (1994) Function of the retinoic acid receptors (RARs) during development. (I) Craniofacial and skeletal abnormalities in RAR double mutants. *Development* **120**, 2723-2748.
- Mackem, S., Knezevic, V. (1999) Do 5'Hoxd genes play a role in initiating or maintaining A-P polarizing signals in the limb? *Cell Tissue Res.* **296**, 27-31.
- Nagy, L., Saydak, M., Shipley, N., Lu, S., Basilion, J. P., Yan, Z. H., Syka, P., Chandraratna, R. A., Stein, J. P., Heyman, R. A., Davies, P. J. (1996) Identification and characterization of a versatile retinoid response element (retinoic acid receptor response element-retinoid X receptor response element) in the mouse tissue transglutaminase gene promoter. *J. Biol. Chem.* **271**, 4355-4365.
- Niederreither, K., Subbarayan, V., Dolle, P., Chambon, P. (1999) Embryonic retinoic acid synthesis is essential for early mouse post-implantation development. *Nat. Genet.* **21**, 444-448.
- Niswander, L., Pizette, S., Zou, H. (2000) BMP function in limb development. *7th International Conference on Limb Development and Regeneration, Aussois, France, 20-24 May 2000*, abstract 10, p. 9.
- Pizette, S., Niswander, L. (2000) BMPs are required at two steps of limb chondrogenesis: formation of prechondrogenic condensations and their differentiation into chondrocytes. *Dev. Biol.* **219**, 237-249.
- Power, S. C., Lancman, J., Smith, S. M. (1999) Retinoic acid is essential for shh/hoxd signaling during rat limb outgrowth but not for limb initiation. *Dev. Dyn.* **216**, 469-480.
- Pravenec, M., Křen, V., Kurtz, T. W., Bílá, V., Kabra, P., Kršiaková, M., Simonet, L. (1990) HXB and BXH sets of recombinant inbred strains: genetic characterization. *Transplant. Proc.* **22**, 2557-2558.
- Robinson, R. (1971) *Gene Mapping of Laboratory Mammals*. Part A. Plenum Press, London-New York.
- Sedmera, D., Novotná, B., Bílá, V., Křen, V. (1998) The role of cell death in limb development of rats manifesting *Lx* allele on different genetic backgrounds. *Eur. J. Morphol.* **36**, 173-181.
- Small, K. M., Potter, S. S. (1993) Homeotic transformations and limb defects in Hox A11 mutant mice. *Genes Dev.* **7**, 2318-2328.
- Stratford, T., Logan, C., Zile, M., Maden, M. (1999) Abnormal anteroposterior and dorsoventral patterning of the limb bud in the absence of retinoids. *Mech. Dev.* **81**, 115-125.
- Sulik, K. K., Dehart, D. B. (1988) Retinoic acid-induced limb malformations resulting from apical ectodermal ridge cell. *Teratology* **37**, 527-537.
- Tsuiki, H., Kishi, K. (1999a) Retinoid-induced limb defects 1: inhibition of cell proliferation in distal mesenchyme of limb buds in rats. *Reprod. Toxicol.* **13**, 103-111.
- Tsuiki, H., Kishi, K. (1999b) Retinoid-induced limb defects 2: involvement of TGF-beta 2 in retinoid-induced inhibition of limb bud development. *Reprod. Toxicol.* **13**, 113-122.
- Underhill, T. M., Weston, A. D. (1998) Retinoids and their receptors in skeletal development. *Microsc. Res. Tech.* **43**, 137-155.
- Vermot, J., Niederreither, K., Dollé, P., Chambon, P. (2000) Initiation of mouse limb bud outgrowth is impaired in the absence of retinoic acid synthesis. *7th International Conference on Limb Development and Regeneration, Aussois, France, 20-24 May 2000*, abstract P4, p. 33.
- Yokouchi, Y., Nakazato, S., Yamamoto, M., Goto, Y., Kameda, T., Iba, H., Kuroiwa, A. (1995) Misexpression of Hoxa-13 induces cartilage homeotic transformation and changes cell adhesiveness in chick limb buds. *Genes Dev.* **9**, 2509-2522.
- Zákány, J., Duboule, D. (1999) Hox genes in digit development and evolution. *Cell Tissue Res.* **296**, 19-25.